



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

GEOMETRY.

343. Proposed by O. J. BROWN, Fairhope, Ala.

From any external point of a triangle, to draw a line so as to divide the triangle into two equal parts.

344. Proposed by C. N. SCHMALL, 604 East 5th Street, New York.

A tinsmith has a sheet of copper in the form of a rectangle, sides a and b . He desires to cut this into two pieces which will form a square when placed together. How can he do this?

CALCULUS.

171. Proposed by E. B. ESCOTT, Ann Arbor, Mich.

In the differential equation

$$\left(\frac{d^2y}{dx^2}\right)^2 \frac{d^5y}{dx^5} - 5\left(\frac{d^2y}{dx^2}\right) \frac{d^3y}{dx^3} \frac{d^4y}{dx^4} + \frac{40}{9} \left(\frac{d^3y}{dx^3}\right)^3 = 0,$$

show that there is an integrating factor of the form $\left(\frac{d^2y}{dx^2}\right)^n$, and integrate the equation.

172. Proposed by CLARENCE OHLENDORF, Chicago, Ill.

Find $\int \log_e \tan^{-1} x dx$.

NUMBER THEORY AND DIOPHANTINE ANALYSIS.

160. Proposed by H. S. VANDIVER, Bala, Pa.

Prove that the integer next above $(1/3+1)^{2n}$ is divisible by 2^{n+1} .

161. Proposed by R. D. CARMICHAEL, Anniston, Ala.

Find a solution of $x^4 - 5x^2 + 4 \equiv 0 \pmod{p \cdot 2p+1}$, where both p and $2p+1$ are odd primes.

162. Proposed by L. E. DICKSON, Ph. D., Associate Professor of Mathematics, The University of Chicago.

If p is an odd prime, find the number of incongruent integers x for which $x^4 + 2ex^2 + f$ is a quadratic residue of p .

MISCELLANEOUS.

181. Proposed by A. H. HOLMES, Brunswick, Me.

In latitude $43^\circ 45'$ N., the sun's declination being $16^\circ 30'$ N., at what time in the forenoon will the angle on the horizon between the east point and the foot of the meridian passing through the sun's position be equal to its altitude?